

LHC PROSPECTS ON HIGGS BOSON SEARCHES

RICCARDO RANIERI

Università degli Studi di Firenze and INFN Firenze

Via G. Sansone, 1

I 50019 Sesto Fiorentino (FI), Italy

e-mail: Riccardo.Ranieri@fi.infn.it

The search for the Higgs boson and physics beyond the Standard Model will be the most important tasks of the two multi-purpose experiments ATLAS and CMS, which will be placed inside the LHC 14 TeV proton-proton accelerator. The most recent studies, which have been developed with detailed simulations of the detector geometry and response, has pointed out that the LHC detectors have the possibility to cover the whole Standard Model Higgs boson mass spectrum and most of the MSSM parameter space.

1. Introduction

The origin of the mass of elementary particles, whose electroweak and strong interactions are described by the Standard Model theory (SM), is thought to be the result of the electroweak symmetry breaking mechanism, which predicts the existence of a new particle, the Higgs boson¹. Experimental observations have confirmed the validity of the Standard Model, but the Higgs boson itself has never been detected. Supersymmetry (SUSY), which relates masses and couplings of scalars and fermions, provides an elegant solution to the *hierarchy problem* of the SM². In the minimal supersymmetric extension of the SM (MSSM)³, each of the known fundamental particles have a super-partner with spin differing by $\frac{1}{2}$ and there are five Higgs scalar mass eigenstates: one CP-odd neutral scalar A, two charged scalars H^\pm and two CP-even neutral scalars h and H (with masses $m_h < m_H$). At tree level the five Higgs bosons masses can be computed in terms of only two parameters, typically chosen to be $\tan\beta$, the ratio between Higgs scalar doublets vacuum expectations values, and m_A , the mass of the pseudoscalar neutral Higgs boson A.

A very powerful accelerator, the Large Hadron Collider (LHC), is being constructed at the European Laboratory of Particle Physics (CERN) in

Geneva, Switzerland, so as to allow a Higgs boson discovery and to search for new physics phenomena up to the TeV energy scale⁴. Two oppositely directed proton beams will be accelerated to a total centre-of-mass energy of 14 TeV and will collide every 25 ns in correspondence of the experimental areas, where particle detectors will be placed. Two of the approved LHC experiments are “A Toroidal LHC ApparatuS” ATLAS⁵ and the “Compact Muon Solenoid” CMS⁶. Both experiments will start data taking during the year 2008, after the LHC machine pilot run scheduled in the Autumn 2007.

2. Standard Model Higgs boson search

The Standard Model is extremely predictive in the Higgs sector, because all couplings, decay widths and production cross sections are given in terms of the unknown Higgs boson mass m_H , being the other parameters experimentally measured. There is not a single production mechanism or decay channel that dominates the whole accessible mass range from 100 GeV/ c^2 to 1 TeV/ c^2 at LHC. Instead several scenarios open up depending on the Higgs boson mass m_H .

The statistical significance for the Standard Model Higgs boson signal observation with 30 fb⁻¹ integrated luminosity is shown in Fig. 1 for m_H ranging between 80 GeV/ c^2 and 1 TeV/ c^2 , for ATLAS⁷ and CMS⁸ experiments. Combining the results for all the channels, almost all the allowed mass range not yet experimentally excluded should be explored during the first year at LHC. After two years about 30 fb⁻¹ integrated luminosity per experiment will be collected and a 7σ signal significance should be reached over the whole mass spectrum, covered by more than one channel.

In the lower mass region, a 2σ significance should be reached during the first year of LHC in different channels: the $t\bar{t}H$ associated production followed by top quarks decays $t \rightarrow bW(\rightarrow \ell\nu)$ ($\ell = e, \mu$) and $t \rightarrow bW(\rightarrow q\bar{q}')$ ⁹, the decay $H \rightarrow \gamma\gamma$ ¹⁰ and the vector boson fusion (VBF) process $qq \rightarrow qqVV \rightarrow qqH$ ($V=W$ or Z)¹¹. The observation of all these channels is very important to extract a convincing signal during the first years of LHC operation. It is likewise important to reduce the uncertainty on background knowledge to less than 10%; this precision could be already reached during the first years of data taking.

At the opening of the ZZ decay channel, the fully leptonic decay $H \rightarrow ZZ^{(*)} \rightarrow 4\ell$ ($\ell = e, \mu$) has the cleanest experimental signature, for this reason it is called the “golden” channel at LHC. The luminosity required for a 5σ discovery is less than 30 fb⁻¹ and corresponds to about two years of data




Figure 1. Expected sensitivity for the observation of the Standard Model Higgs boson signal as a function of the mass m_H with ATLAS (a) and CMS (b) with 30 fb^{-1} integrated luminosity.

taking at low luminosity¹². If $m_H < 2m_Z$ only one Z can be reconstructed. As a result higher luminosity is needed to observe the signal.

3. MSSM Higgs boson search

3.1. Neutral Higgs bosons h, H and A

The dominant production mechanism in the phenomenologically relevant Higgs mass ranges for small and moderate values of $\tan\beta$ is the gluon-gluon fusion process. At large $\tan\beta$, the associated production involving a $b\bar{b}$ pair is enhanced by the large Yukawa couplings to down-type quarks. The main decays are into $b\bar{b}$ (about 90%) and $\tau^+\tau^-$ pairs (about 10%). A clean signature can be extracted by the $h/H/A \rightarrow \mu^+\mu^-$ decay channel, although the branching ratio into $\mu^+\mu^-$ is 3×10^{-4} over all the mass range for $\tan\beta=30$. Nevertheless there are some factors that can provide an observable rate and make this channel appealing: at large $\tan\beta$ rates are enhanced through associated $b\bar{b}h/H/A$ production and, to some extent, the reduction in signal rate with respect to the $\tau^+\tau^-$ channel is compensated by the much better identification efficiency achieved in the $\mu^+\mu^-$ channel. If the neutral Higgs boson mass is not too close to the Z mass, with 30 fb^{-1} of integrated luminosity the signal significance is greater than 5σ and thus a discovery for $\tan\beta=30$ is possible¹³.

3.2. Charged Higgs Bosons

The search for the MSSM charged Higgs boson H^\pm is performed in two different ways depending on the mass m_{H^\pm} and the dominant decay channels. The light charged Higgs boson is searched in the $H^\pm \rightarrow \tau\nu_\tau$ decay channel, whose branching ratio is about 100%. It is produced by top quark decays $t \rightarrow bH^\pm$. The background with an isolated lepton from the accompanying W decay in $t \rightarrow bW^\pm$ is suppressed by searching for channels with a single τ jet or two τ jets in the final state. A 5σ significance can be reached¹⁴ with 30 fb^{-1} if $m_{H^\pm} < 100 \text{ GeV}/c^2$.

The heavy charged Higgs boson is searched for exploiting the main decay channel $H^\pm \rightarrow tb$ in the associated production process $gg \rightarrow btH^\pm$, even with low signal-to-background ratios due to the $t\bar{t}$ multi-jet background. Also the $H^\pm \rightarrow \tau\nu_\tau$ and $H^\pm \rightarrow Wh$ decays can be studied. The $H^\pm \rightarrow \pi\nu_\tau$ is particularly interesting when hadronic τ decays are required, because the $t\bar{t}$ and tW backgrounds can be suppressed exploiting the different helicity correlations in the $H^\pm \rightarrow \pi\nu_\tau$ and $W^\pm \rightarrow \pi\nu_\tau$ decays. The $H^\pm \rightarrow \mu\nu_\mu$ decay can be efficiently extracted with integrated luminosity exceeding 100 fb^{-1} .

References

1. P.W. Higgs, *Phys. Lett.*, **12** (1964) 132-133, *Phys. Rev. Lett.*, **13** (1964) 508-509, *Phys. Rev.*, **145** (1966) 1156-1163; F. Englert and R. Brout, *Phys. Rev. Lett.*, **13** (1964) 321-322; G.S. Guralnik, C.R. Hagen and T.W.B. Kibble, *Phys. Rev. Lett.*, **13** (1964) 585-587.
2. P. Fayet, *Phys. Lett.*, **B64** (1976) 159, *Phys. Lett.*, **B69** (1977) 489; G. R. Farrar and P. Fayet, *Phys. Lett.*, **B76** (1978) 575-579.
3. H. P. Nilles, *Phys. Rept.*, **110** (1984) 1; H. E. Haber, *Phys. Rept.*, **117** (1985) 75; L. E. Ibanez, G. G. Ross, *Phys. Lett.*, **B110** (1982) 215-220; S. Dawson, hep-ph 9712464.
4. LHC Study Group, CERN/AC 95-05.
5. ATLAS Collaboration, CERN/LHCC 94-43, LHCC/P2.
6. CMS Collaboration, CERN/LHCC 94-38, LHCC/P1.
7. ATLAS Collaboration, CERN/LHCC 99-15 ATLAS TDR 15.
8. S. Abdullin *et al.*, CMS NOTE 2003/033.
9. J. Camin and M. Schumacher, ATL-PHYS-2003-024.
10. ATLAS Collaboration, CERN/LHCC 96-40 ATLAS TDR 1; CERN/LHCC 96-41 ATLAS TDR 2.
11. S. Asai *et al.*, arXiv:hep-ph/0402254.
12. M. Sani, *Czech. J. Phys.* **55** (2005) B101.
13. S. Marcellini and G. Masetti, CMS NOTE in preparation.
14. R. Kinnunen, CMS NOTE-2006/100; M. Baarmand, M. Hashemi and A. Nikitenko, CMS NOTE-2006/056.